

PROF. XIN-LI XIA (Orcid ID : 0000-0003-3731-4970)

Article type : Research Article

PeCHYR1, a ubiquitin E3 ligase from *Populus euphratica*, enhances drought tolerance via ABA-induced stomatal closure by ROS production in *Populus*

Fang He, Hou-Ling Wang, Hui-Guang Li, Yanyan Su, Shuang Li, Yanli Yang, Cong-Hua Feng,

Weilun Yin, Xinli Xia*

Beijing Advanced Innovation Center for Tree Breeding by Molecular Design, National Engineering

Laboratory for Tree Breeding, College of Biological Sciences and Technology, Beijing Forestry

University, Beijing 100083, P.R. China

* Corresponding author: Xinli Xia (Tel : 008610-62336400; Fax: 8610-62336400; E-mail addresses:

xiaxl@bjfu.edu.cn

Running title: *PeCHYR1* overexpression enhances drought tolerance.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/pbi.12893

This article is protected by copyright. All rights reserved.

Key words: drought tolerance; stomatal closure; abscisic acid; *PeCHYRI*; *Populus*.

Summary

Drought, a primary abiotic stress, seriously affects plant growth and productivity. Stomata play a vital role in regulating gas exchange and drought adaptation. However, limited knowledge exists of the molecular mechanisms underlying stomatal movement in trees. Here, *PeCHYRI*, a ubiquitin E3 ligase, was isolated from *Populus euphratica*, a model of stress adaptation in forest trees. *PeCHYRI* was preferentially expressed in young leaves and was significantly induced by ABA (abscisic acid) and dehydration treatments. To study the potential biological functions of *PeCHYRI*, transgenic poplar 84K (*Populus alba* × *Populus glandulosa*) plants overexpressing *PeCHYRI* were generated. *PeCHYRI* overexpression significantly enhanced H₂O₂ production and reduced stomatal aperture. Transgenic lines exhibited increased sensitivity to exogenous ABA and greater drought tolerance than that of WT (wild-type) controls. Moreover, up-regulation of *PeCHYRI* promoted stomatal closure and decreased transpiration, resulting in strongly elevated WUE (water use efficiency). When exposed to drought stress, transgenic poplar maintained higher photosynthetic activity and biomass accumulation. Taken together, these results suggest that *PeCHYRI* plays a crucial role in enhancing drought tolerance via ABA-induced stomatal closure caused by hydrogen peroxide (H₂O₂) production in transgenic poplar plants.

Introduction

Drought stress is a common abiotic stress, affecting plant growth, seed production, gas exchange, water relations and cellular homeostasis in plants (Shinozaki and Yamaguchi-Shinozaki, 2007; Zhu, 2016). To adapt to hostile environments, plants have developed intrinsic mechanisms to mitigate drought stress, such as closing stomata, reducing transpiration, generating abscisic acid (ABA), and accumulating hydrogen peroxide (H_2O_2) (Dietz et al., 2016; Zhu, 2016). ABA-induced stomatal closure is a defensive action that is usually accompanied by the production of H_2O_2 (Li et al., 2017; Zhang et al., 2001). NADPH oxidase (RbohF) can generate H_2O_2 in guard cells, which crucially modulates ABA-induced stomatal closure and plant drought tolerance (Gudesblat et al., 2007; Kolla et al., 2007; Ma et al., 2016). Previous studies have indicated that AtRbohF can be phosphorylated by OST1 (an ABA-induced SnRK2.6 protein kinase) (Joshi-Saha et al., 2011; Sirichandra et al., 2009; Wege et al., 2014). H_2O_2 is a reactive oxygen species (ROS) that acts as a major signaling molecule in the ABA response pathway in guard cells (Bright et al., 2006; Li et al., 2017). High concentrations of ROS lead to cell injury or even hypersensitive cell death, whereas low concentrations of ROS function as developmental signals, controlling all aspects of plant biology (Ahmad et al., 2008; Dietz et al., 2016; Huang et al., 2016; Karuppanapandian et al., 2011; Pitzschke et al., 2006). OsASR5 and DCA1, which enhance drought tolerance in rice, control stomatal closure by adjusting the concentration of H_2O_2 (Cui et al., 2015; Li et al., 2017). GHR1 modulates stomatal movement associated with H_2O_2 signaling in Arabidopsis (Hua et al., 2012).

In stomata, ABA-mediated ROS generation elevates cytosolic calcium ion levels and gives rise to stomatal closure (McAinsh et al., 1996; Pei et al., 2000; Xing et al., 2008). Recently, it was reported that ROS can induce stomatal closure through an ABA-independent pathway in rice (*Oryza sativa*).

This article is protected by copyright. All rights reserved.

The SNAC1-targeted gene OsSRO1c positively controls H₂O₂-induced stomatal closure by regulating hydrogen peroxide level in rice, and the E3 ligase OsHTAS can promote heat tolerance by modulating H₂O₂-induced stomatal closure in rice (Liu et al., 2016; You et al., 2013). However, 35S:OsASR5 plants were more sensitive to exogenous ABA treatment, activating the production of H₂O₂ (Li et al., 2017).

RING (really interesting new gene) finger proteins consist mainly of conserved “CxHY” domains (McDowall, 2007). These conserved “CxHY” motifs are part of the CHY zinc-finger domain, which plays a role in physical interaction and ubiquitination (Lee and Kim, 2011; Wooff et al., 2004; Yu et al., 2016), contains 12 histidines and cysteines, and binds with three zinc ions, composing a unique zinc-finger motif (Cayrol et al., 2007; Ding et al., 2015b; Esposito et al., 2006). Proteins such as ubiquitin (ubi) E3 ligases, a core component of ubiquitination pathway containing a CHY zinc-finger domain, possess a universal function including ubiquitylation of substrate proteins (Lim et al., 2013a; Ning et al., 2016; Stone et al., 2005). The ubiquitination process plays a crucial role in plant development and response to environmental stresses (Lim et al., 2013a; Stone et al., 2005). RING ubi E3 ligases play decisive roles in almost all plant growth processes, including photomorphogenesis, flower development, phytohormone signaling, and regulation of senescence (Bae et al., 2011; Ding et al., 2015b; Lim et al., 2017; Lim et al., 2013b). RING E3 ligases have been found in diverse plants, such as hot pepper (*Capsicum annuum*), maize (*Zea mays*), Arabidopsis and rice, positively or negatively regulating abiotic stress (Lyzena and Stone, 2012; Park et al., 2015; Zhao et al., 2014). Recently, a ubiquitin E3 ligase, Arabidopsis CHYR1, was found to positively facilitate ABA and drought-mediated stomatal closure, ROS production, and plant drought tolerance via

This article is protected by copyright. All rights reserved.

Accepted Article

SnRK2.6-mediated phosphorylation (Ding et al., 2015a). However, the expression of a homolog of *CHYR1*, rice *OsRZFP34*, enhances stomatal opening, leaf cooling, and ABA insensitivity (Hsu et al., 2014a).

Although the roles of CHYR1 family proteins have been widely reported in herbaceous plants, their possible functions in ligneous plants, especially in stomatal movement, remain unknown. Several mechanisms of plant response to drought exist, including avoidance, escape and tolerance strategies. Tolerance mechanisms attempt to maintain plant functions at the same level as in unstressed conditions, while avoidance mechanisms are used to adjust the balance between water loss and water uptake (Moradi, 2016). *Populus*, a pioneer genus in forest ecosystems, possesses great economic and ecological value (Sterky et al., 2004). Most poplar plants are fast-growing and have poor drought tolerance (Larcheveque et al., 2011; Tuskan et al., 2006; Windt et al., 2006). However, one exception, *P. euphratica*, is the only large tree species that forms forests in desert areas, playing a very important role in the maintenance of the local ecological balance. Due to its strong adaptability to extreme temperature, drought and salinity, *P. euphratica* has served as a model woody plant to explore the mechanisms of stress resistance (Chen et al., 2013; Li et al., 2011; Ma et al., 2013; Tang et al., 2013; Yan et al., 2012). From an analysis of a transcriptome from *P. euphratica* under drought stress, we selected *PeCHYR1*, which is up-regulated in drought-stressed *P. euphratica* leaves, and performed further functional analysis of this gene in poplar (Tang et al., 2013). In the current study, *PeCHYR1* was cloned from *P. euphratica* and transferred into poplar 84K (*P. alba* × *P. glandulosa*) (Feng et al., 2015; Ke et al., 2016). Next, we verified the function of *PeCHYR1* by means of molecular biology and plant physiological indexes. We found that *PeCHYR1* was a functional homolog of *CHYR1*.

Overexpression of *PeCHYR1* enhanced drought tolerance by promoting H₂O₂-mediated stomatal closure in poplar.

Results

Isolation and sequence analysis of *PeCHYR1*

To elucidate the potential function of *PeCHYR1* in the abiotic stress response of ligneous plants, we took advantage of *P. euphratica*, which naturally grows in deserts, to clone *PeCHYR1*. The coding sequence length of *PeCHYR1* was 876 bp, encoding 291 amino acids. Homology analyses of the amino acid sequence showed 52 homologous genes in *Populus*, *Amborella*, *Eucalyptus*, *Malus*, *Prunus*, *Theobroma*, *Picea*, *Pinus*, *Salix*, *Arabidopsis*, rice, and maize. The evolutionary tree of related genes was divided into four parts (I, II, III, and IV). The nucleic acid sequence of *PeCHYR1* is mostly identical to that of *PtrCHYR1* (Potri.009G005700), sharing 96.7% sequence identity (Figure 1a). A multiple sequence alignment revealed that the amino acid sequences of *PeCHYR1*, *PtrCHYR1*, *SpCHYR1* and *AtCHYR1* (AT5G22920) contained the same conserved domains (Figure 1b). Furthermore, the *PeCHYR1* promoter contains CAAT and TATA motifs that are involved in drought stress response (Figure S1).

PeCHYR1 involvement in ABA and water stress response

To study the potential biological functions of *PeCHYR1*, we applied quantitative real-time polymerase chain reaction (RT-qPCR) analysis to determine the relative transcript abundance in various tissues of

P. euphratica. We observed that *PeCHYR1* was mainly expressed in the leaf rather than the other tissues (Figure 2a). *P. euphratica* plants were treated with exogenous ABA and drought stress to investigate the expression pattern of *PeCHYR1*. The results showed that the transcript abundance of *PeCHYR1* is transiently increased in the leaf under water stress, reaching a maximum of 3.5 times higher than control at 3 h, dropping to approximately 3 times at 6 and 9 hours, and decreasing to 1.2 times at 12 hours (Figure 2b). The abundance of *PeCHYR1* was gradually augmented by ABA treatment, significantly increasing at 0.5 h, maintaining a stable level until 6 h, and reaching a maximum of 9 times higher than control at 12 h (Figure 2c).

Drought-stress-induced ABA generation leads to the accumulation of ROS. Young leaves of 1.5-month-old poplar were treated with 100 μ M ABA for 0, 1, 2, and 3 h, and then the leaves were immersed in diaminobenzidine (DAB) overnight. We observed that the red-brown color of the leaves darkened gradually as the time of ABA treatment progressed (Figure 2d).

Subcellular localization of PeCHYR1

Previous studies indicated that Arabidopsis CHYR1, a ubiquitin E3 ligase, was generally localized in the nucleus, cytoplasm, and endoplasmic reticulum (ER) (Ding et al., 2015b). Thus, *P. euphratica* PeCHYR1, a functional homolog of CHYR1, might localize to the same cellular structures. To determine the subcellular localization of PeCHYR1, a 35S:PeCHYR1-GFP (green fluorescent protein) fusion protein, together with a 35S: HDEL-RFP (red fluorescent protein) fusion protein, was transiently transfected into tobacco leaves and Arabidopsis leaf protoplasts. Co-localization of PeCHYR1-GFP and HDEL-RFP was distinctly detected in the ER (Figure 3a, b). Simultaneously,

This article is protected by copyright. All rights reserved.

co-localization with 4', 6-diamidino-2-phenylindole (DAPI) dye proved that the 35S:PeCHYR1-GFP fusion protein was localized to the nucleus (Figure 3a, b). Overall, the results indicated that PeCHYR1 was localized to the nucleus and ER.

To study the potential biological functions of *PeCHYR1*, transgenic poplar 84K (*P. alba* × *P. glandulosa*) plants overexpressing *PeCHYR1* were generated. Every transgenic line was verified by PCR, RT-qPCR, and histochemical staining with β -glucuronidase (GUS) (Figure S2). In the literature, these three methods are usually used to verify transgenic plants (Jin et al., 2017).

***PeCHYR1* promotes ABA-induced stomatal closure via ROS production**

ABA and ROS can induce stomatal closure (Bright et al., 2006), so we considered whether *PeCHYR1* could regulate ABA-induced ROS accumulation. We treated 35S:*PeCHYR1* (*OXPeCHYR1-1*, *OXPeCHYR1-7*, *OXPeCHYR1-8*) and WT plants with 100 μ M ABA (0, 1 and 3 h) and then performed DAB staining on their leaves. When the leaves of the different lines were treated with ABA, we found that the brown color of 35S:*PeCHYR1* leaves was deeper than that of WT leaves (Figure 4a), even after multiple hours of treatment. These results indicated that 35S:*PeCHYR1* plants had higher ROS levels than WT plants in the instantaneous ABA treatment.

To confirm whether *PeCHYR1* participated in ABA-induced H₂O₂ signaling, we measured the endogenous H₂O₂ levels in the stomata of WT and 35S:*PeCHYR1* poplars. We used 2,7-dichlorodihydrofluorescein diacetate (H₂DCF-DA), a unique fluorescent probe, to detect the H₂O₂ levels within guard cells. In the absence of ABA, we found that 35S:*PeCHYR1* plants showed slightly elevated H₂O₂ content compared with WT plants (Figure 4b, c). In the presence of ABA, the levels of

This article is protected by copyright. All rights reserved.

H₂O₂ were increased in both WT and 35S:*PeCHYRI* poplar, while 35S:*PeCHYRI* poplar exhibited significantly higher H₂O₂ levels than WT plants (Figure 4b, c).

To explore whether *PeCHYRI* promotes ABA-induced stomatal closure, stomatal movement in response to ABA treatment was observed in WT and 35S:*PeCHYRI*. Then, leaf stomata were observed by scanning electron microscopy. Meanwhile, we observed a significant difference when the leaves were treated with 5 μM ABA for 1 and 2 hours; the stomatal apertures closed more quickly in the leaves of the transgenic plants than in leaves of the WT plants (Figure 4d, e). Conclusively, these data indicated that *PeCHYRI* actively regulated ABA-induced ROS production.

***PeCHYRI* overexpression enhances WUE by reducing stomatal conductance**

As H₂O₂ can also induce stomatal closure (Bright et al., 2006), we hypothesized that overexpression of *PeCHYRI* in transgenic poplar plants may decrease stomatal opening. To determine whether *PeCHYRI* could affect photosynthesis in plants, we measured photosynthesis-light curves in WT and 35S:*PeCHYRI* poplars. The data indicated that the patterns of photosynthesis were roughly similar in WT and 35S:*PeCHYRI* poplars (Figure 5a). The stomatal conductance (Gs) data indicated that WT and 35S:*PeCHYRI* poplars had altered Gs responses to light, while the Gs of 35S:*PeCHYRI* poplars was lower than that of WT (Figure 5b).

At the same time, leaf transpiration in 35S:*PeCHYRI* poplars was significantly lower than that in WT poplars (Figure 5b). In general, instantaneous WUE values in 35S:*PeCHYRI* plants were higher than those in WT plants (Figure 5d). Vapor pressure deficit (VPD) was measured, and no significant

Accepted Article

differences were found between WT and 35S:*PeCHYR1* poplars (Figure 5e). This lack of a significant difference in VPD indicates that the differences in transpiration rate were not caused by VPD.

35S:*PeCHYR1* poplars possess drought tolerance

Because ABA and drought stress can induce the expression of *PeCHYR1* (Figure 2b, c), we hypothesized that *PeCHYR1* played a vital role in drought response. Afterwards, 35S:*PeCHYR1* poplars and WT poplars were cultured under the same conditions and then subjected to a drought treatment in which the soil relative water content (RWC) was reduced from 70% and rewatered again for 2 days. On day 3, most leaves of the WT poplars were slightly wilted, while the transgenic poplars still appeared normal. On day 7, the leaves of the WT poplars were seriously wilted, while those of the transgenic poplars remained turgid. Furthermore, when the plants were watered again after drought stress, the WT poplars were unable to recover completely, whereas the transgenic poplars returned to normal and continued to grow (Figure 6a). Control plants were kept under the same conditions, except that the soil RWC was maintained at 70%, and no significant difference in physiological indicators and phenotype was found between 35S:*PeCHYR1* poplars and WT poplars in control conditions (Figure 6).

Leaf RWC reflects the degree of water deficit imposed. When exposed to short-term drought stress, 35S:*PeCHYR1* poplar retained observably higher leaf RWC values than WT poplar (Figure 6b). As malondialdehyde (MDA) content is a valid indicator of cytomembrane oxidative damage, we measured the variations in MDA content caused by drought stress in the leaves of WT and transgenic. This article is protected by copyright. All rights reserved.

Accepted Article

poplars. The results indicated that the 35S:*PeCHYR1* poplars had lower levels of MDA than the WT poplars did when exposed to short-term drought stress (Figure 6c). The REC (relative electrical conductance) of both 35S:*PeCHYR1* and WT poplars increased after drought treatment. After drought stress, the transgenic poplars showed less membrane damage (approximately 24.8%–27.9% ion leakage), whereas the leaves of WT poplars showed severe membrane damage (approximately 57.6% ion leakage) (Figure 6d).

To study the effects of drought stress in regard to H₂O₂, we measured H₂O₂ contents after 0, 3 and 7 days of drought stress. After withholding water for 0 days, the H₂O₂ contents in transgenic poplars were slightly higher than those in WT plants. The 35S:*PeCHYR1* transgenic and WT poplars both had increased H₂O₂ levels after 3 days and 7 days of drought stress; however, the H₂O₂ content in WT poplars was significantly higher than that in transgenic poplars (Figure 6e). Peroxidase (POD) and superoxide dismutase (SOD) are crucial antioxidants that can scavenge ROS. Thus, we detected the activities of these two enzymes in 35S:*PeCHYR1* and WT poplars before and after drought treatment. Corresponded with the H₂O₂ levels, the results indicated that the activity levels of POD and SOD were dramatically elevated in 35S:*PeCHYR1* plants in comparison to WT plants under drought stress conditions (Figure 6f, g)

As shown in Figure 7a–c, the 35S:*PeCHYR1* transgenic and WT poplars during the 6 days of drought stress had remarkable variations in photosynthesis, Gs, and transpiration. During these 6 days, photosynthesis in the transgenic and wild-type poplars both decreased, while photosynthesis in WT poplars decreased more rapidly than that in the transgenic poplars under drought conditions, and

Accepted Article

photosynthesis in WT poplars was lower than that in the transgenic poplars after 2 days of drought treatment (Figure 7c). As with Gs, transpiration in the transgenic and WT poplars showed an overall decreasing trend, while transpiration in WT poplars decreased faster than that in transgenic poplars under drought conditions, and transpiration in WT poplars was lower than that in transgenic poplars after 3 days of drought stress (Figure 7a, b). Moreover, the data indicated that isolated leaves from WT poplars lose water faster than those from transgenic poplars under natural dehydration (Figure 7d). Conclusively, 35S:*PeCHYRI* transgenic poplars had better tolerance to drought stress.

35S:*PeCHYRI* poplars exhibited enhanced drought tolerance under long-term water stress

To thoroughly study how *PeCHYRI* expression in plants influences the response to long-term water deficit, 35S:*PeCHYRI* poplars and WT poplars were subjected to a 32-day water deficit assay by holding the soil RWC at stable values (45% and 70%). After 32 days, in both shoot and root, the conditions of the WT and 35S:*PeCHYRI* poplar were approximately identical at 70% soil RWC. After 32 days of drought treatment, 35S:*PeCHYRI* poplar reached greater heights and appeared in better condition than WT at 45% soil RWC (Figure 8a). Furthermore, at 70% soil RWC, the chlorophyll level of 35S:*PeCHYRI* poplar was similar to that of WT plants. However, after the 32-day water deficit assay at 45% soil RWC, 35S:*PeCHYRI* poplar had higher chlorophyll content than WT plants (Figure 8b). In addition, 35S:*PeCHYRI* poplar showed higher photosystem II efficiency (Fv/Fm) and higher leaf RWC than WT poplar did (Figure 8c, d).

To explore whether *PeCHYRI* would affect growth and development in the absence of water stress (70% soil RWC) and under mild water stress (45% soil RWC), plant height and stem height growth

Accepted Article

rate were monitored. The 35S:*PeCHYRI* poplar had similar plant height and stem height growth rate to those of wild-type plants in the absence of water stress (Figure 9a, b). At 45% soil RWC, the 35S:*PeCHYRI* poplar had a significantly greater plant height and stem height growth rate than those of WT plants (Figure 9a, b). These results suggested that 35S:*PeCHYRI* poplar maintained higher growth under long-term water deficit conditions.

To observe the difference in biomass accumulation between WT and 35S:*PeCHYRI* poplar under no water stress (70% soil RWC) and mild water stress (45% soil RWC), shoot weight, root weight, total biomass, and root-shoot ratio were monitored every eight days. Under no water stress, transgenic and WT plants had similar biomass accumulations in all parts of the plant (Figure 9c–f). However, compared with that of WT plants, transgenic plants had more biomass accumulation in the shoot and root under mild water stress, and the shoot weight, root weight, and total biomass of 35S:*PeCHYRI* poplar were obviously higher than those of WT plants after 8 days of drought treatment (Figure 9c–e). The root-shoot ratio reflects the relationship between the aboveground and underground parts of the plant. Both 35S:*PeCHYRI* and WT plants had greater root-shoot ratios, and 35S:*PeCHYRI* poplar had a larger root-shoot ratio than that of WT after 16 days of drought treatment (Figure 9f). After 32 days of drought treatment, the biomass of 35S:*PeCHYRI* poplar was 36.9 and 43.4% higher, respectively, than that of WT (Figure 9e), and the root-shoot ratio of 35S:*PeCHYRI* poplar was 15.3% and 8.9% higher, respectively, than that of WT (Figure 9f). In summary, under water deficit conditions, 35S:*PeCHYRI* poplar maintained higher growth than the WT.

***PeCHYR1* mediated the expression of upstream and downstream genes involved in adversity**

The genes *LEA14*, *RbohD*, *RbohF* and *Snrk2.6* are known to be drought responsive and have positive functions in drought tolerance (Ding et al., 2015a). Thus, WT and 35S:*PeCHYR1* poplars were treated with 2 h dehydration stress to investigate changes in *PeLEA14*, *PeRbohD*, *PeRbohF* and *PeSnrk2.6* expression. The results showed that the expression levels of *PeLEA14*, *PeRbohD*, *PeRbohF* and *PeSnrk2.6* in transgenic poplars were greater than those in WT poplars after 2 h dehydration stress (Figure 10a–d). These results indicated that the improved drought tolerance of 35S:*PeCHYR1* transgenic poplars may depend on the regulation of downstream genes.

Discussion

PeCHYR1 was cloned from *P. euphratica*. A multiple alignment of amino acid sequences showed that *PeCHYR1*, *PtRCHYR1*, *SpCHYR1* and *AtCHYR1* all contain a RING domain and a CHY zinc-finger domain (Figure 1b). The conserved “CxHY” motif lies in the CHY zinc-finger domain, which plays a role in physiological interaction and ubiquitination (Lee and Kim, 2011; Leng et al., 2003) and consists of 12 histidines and cysteines that bind with three zinc ions, composing a unique zinc-finger motif (Cayrol et al., 2007). Phylogenetic analysis indicated that the sequences of *PeCHYR1*, *PtRCHYR1*, and *AtCHYR1* had higher homology (Figure 1a), potentially sharing a conserved and crucial function. Based on the evolutionary tree of related genes, *PeCHYR1*, *PtCHYR1* (Potri.009G005700), and *AtCHYR1* (AT5G22920.1) belonged to the first part (I) in the same branch (Figure 1a), which may decrease relative stomatal aperture and enhance drought tolerance under stress control conditions (Ding et al., 2015b). However, *OsRZFP34*, *AtCHYR2* (AT5G25560.2), and

This article is protected by copyright. All rights reserved.

PtCHYR2 (Potri.006G245400) belonged to the third part (III) in the same branch (Figure 1a), which may control stomatal opening even with ABA treatment (Hsu et al., 2014a). *PeCHYR1* also possesses a particular function; this protein is probably phosphorylated by *PeSnRK2.6* and augments drought and ABA-induced responses in plants, which is similar to the function of *CHYR1* in Arabidopsis (Ding et al., 2015a). Furthermore, transgenic poplar plants overexpressing *PeCHYR1* showed increased drought tolerance via ABA-induced stomatal closure and ROS production.

Previous studies have indicated that plants regulate ABA-induced stomatal closure by ROS accumulation (Bright et al., 2006; Hua et al., 2012; Li et al., 2017). H_2O_2 is a ROS that acts as a second messenger in ABA signaling in guard cells (Bright et al., 2006; Dietz et al., 2016; Kohler et al., 2003). The content of H_2O_2 in plants increases considerably under treatment with exogenous ABA (Figure 2d). This increased H_2O_2 level is an early signaling event in ABA-induced signal transduction (Bright et al., 2006; Dietz et al., 2016; Kreslavski et al., 2012). In addition, the level of H_2O_2 increased more under treatment with exogenous ABA in 35S:*PeCHYR1* transgenic plants than in WT plants (Figure 4a–c). The altered relative expression levels of *PeRbohD* and *PeRbohF* (Figure 10c, d), which are involved in producing H_2O_2 , were in agreement with the above results. Overexpression of *PeCHYR1* augmented ABA responsiveness, thereby activating stress-inducible gene expression.

These results indicated that *PeCHYR1* may promote ABA-induced H_2O_2 production.

ABA is involved in abiotic stress response and regulates the expression of many stress-responsive genes (Danquah et al., 2014; Kobayashi et al., 2005; Zhu, 2016). Whether these genes are induced in response to exogenous ABA treatment depends on their involvement in the ABA-independent or ABA-dependent stress response pathways (Yang et al., 2011). The protein DST generates an increase of hydrogen peroxide and promotes stomatal closure by an ABA-independent pathway (Huang et al., 2011). This article is protected by copyright. All rights reserved.

2009). However, 35S:*OsASR5* plants were more sensitive to exogenous ABA treatment than WT plants, suggesting that *OsASR5* was involved in an ABA-dependent pathway (Li et al., 2017). In our study, treatment with exogenous ABA did observably affect the expression of *PeCHYRI* (Figure 2c), and significant differences were present between the stomatal apertures of the WT and 35S:*PeCHYRI* lines under ABA treatment (Figure 4d, e), demonstrating that *PeCHYRI* increased sensitivity to exogenous ABA and reduced stomatal aperture. At the same time, *PeCHYRI* also participates in biological stress responses (Figure S3a, b).

Stomata control gas exchange, and they can regulate transpiration by limiting water loss and affect photosynthesis by facilitating CO₂ uptake (Hetherington and Woodward, 2003; Vahisalu et al., 2008).

Stomatal conductance is a direct result of stomatal density and aperture (Bussis et al., 2006; Chaerle et al., 2005; Qin et al., 2016). In our study, changes in transpiration were consistent with changes in stomatal conductance. As same as stomatal conductance, transpiration in 35S:*PeCHYRI* plants was significantly lower than that in WT plants (Figure 5b, c). However, the change of photosynthesis in WT plants was roughly similar to that in 35S:*PeCHYRI* plants. Accordingly, the instantaneous WUE in the transgenic plants was elevated (Figure 5d), consistent with the normal-temperature water loss rate of 35S:*PeCHYRI* plants, which showed less water loss than WT plants did (Figure 7d). Previous studies have indicated that the decreased stomatal conductance in 35S:*PdEPF1* plants would not necessarily lead to decreased photosynthesis under well-watered conditions (Wang et al., 2016).

Rubisco and ribulose-1,5-bisphosphate are nonstomatal factors that play roles in modulating photosynthetic rate (Percy and Seemann, 1990; Qin et al., 2016), and thus, stomatal conductance does not solely control changes in foliar photosynthetic rate (Niinemets, 2002). These results indicate

that a modest decrease in stomatal conductance in 35S:*PeCHYRI* transgenic plants can increase WUE but not affect photosynthesis under unstressed conditions.

When the plants were in a strictly water-controlled environment, the reduced stomatal conductance of the 35S:*PeCHYRI* transgenic plants possibly reduced water evaporation and played an important part in maintaining a higher leaf RWC (Figure 7b). Under ABA and dehydration treatment, the expression level of *PeCHYRI* gradually rose (Figure 2b, c), consistent with the characteristics of drought tolerance in 35S:*PeCHYRI* transgenic plants. The evidence indicated that 35S:*PeCHYRI* plants had a stronger drought tolerance phenotype than WT plants did. Under drought conditions, 35S:*PeCHYRI* transgenic plants could assimilate more carbon dioxide (Figure 7c). Generally, the level of H₂O₂ in plants can significantly increase under stress conditions (Bhattacharjee, 2012; Bright et al., 2006; de la Garma et al., 2015). Abrupt increases in plant H₂O₂ content are regarded as a marker of various stresses that can induce the antioxidant defense system (Ding et al., 2010; Jiang and Zhang, 2002). Nevertheless, exceedingly high levels of H₂O₂ can give rise to oxidative injury (Kuge et al., 2010; Wang et al., 2008). In our study, both in the short-term ABA treatment and early in the drought treatment, the endogenous hydrogen peroxide levels in 35S:*PeCHYRI* transgenic plants suddenly increased and became higher than those in the WT plants (Figure 4a–c), thereby activating the native antioxidant system to cope with the stress. However, when the plants were in extreme drought conditions, the accumulation of H₂O₂ in the WT was significantly higher than that in the overexpression lines (Figure 6e), the stomatal conductance in the WT lines almost reached zero (Figure 7b), and photosynthesis was markedly lower than that in the overexpression lines (Figure 7c). The major ROS-scavenging mechanisms include APX (ascorbate peroxidase), POD, SOD, and CAT (catalase) (Ahmad et al., 2008; Dietz et al., 2016; Willekens H, 1997), and 35S:*PeCHYRI* plants

This article is protected by copyright. All rights reserved.

showed higher activity levels of POD and SOD under drought stress conditions. POD and SOD, which are beneficial to the maintenance of ROS levels during long-term drought, were activated in transgenic plants (Figure 6f, g). Our observations demonstrated that 35S:*PeCHYRI* transgenic plants underwent a sudden, dramatic increase of H₂O₂, which activated the antioxidant system to reduce oxidative damage after drought treatment.

In addition, the level of MDA indirectly indicates the extent of membrane lipid peroxidation (Shi et al., 2013b; Shi et al., 2014b). In our research, drought stress led to mass accumulation of ROS in WT plants, elevating the content of MDA compared with that in 35S:*PeCHYRI* transgenic plants, seriously damaging membrane permeability, causing intracellular ion leakage, and thereby exacerbating plant senescence and death (Figure 6c, d). Thus, WT plants continued to receive more damage than transgenic plants, and we conclude that overexpression of *PeCHYRI* in plants is beneficial for drought tolerance.

Photosystem II efficiency (Fv/Fm), which reflects the potential maximum light energy conversion efficiency of plants, is 0.8–0.85 under normal conditions in the vast majority of higher plants (Kitajima and Butler, 1975). Photosystem II efficiency declines in plants under environmental stress (Ke et al., 2016; Winter and Lesch, 1992). Although drought stress affects plant photosystem II electron transport and reduces chlorophyll content, 35S:*PeCHYRI* poplar exhibited higher chlorophyll content and photosystem II efficiency than WT poplar did (Figure 8b, c), leading to less injury under water deficit conditions (Figure 8a). Furthermore, we found that leaf RWC in 35S:*PeCHYRI* poplar was obviously higher than that in WT poplar (Figure 8d), which also ensured that plant photosystem II could transport electrons properly and that plants could assimilate carbon dioxide normally under severe drought stress. Our experiment clearly demonstrated that poplars with decreased stomatal

This article is protected by copyright. All rights reserved.

conductance could increase in plant height and biomass under water deficit conditions (Figure 9).

35S:*PeCHYR1* poplar showed decreased transpiration, increased photosynthetic rate, enhanced WUE, elevated plant height, and boosted biomass under water deficit conditions. Drought is the primary abiotic stress responsible for negatively influencing poplar growth (Dash et al., 2017; Ke et al., 2016; Yin et al., 2004). We suggest that 35S:*PeCHYR1* poplar, a novel and drought-tolerant forestry species, can improve WUE and increase biomass under drought stress.

In summary, a ubiquitin E3 ligase from *P. euphratica*, *PeCHYR1*, increased sensitivity to exogenous ABA, enhanced ROS production and reduced stomatal aperture, thereby reinforcing plant drought tolerance. Moreover, the up-regulation of *PeCHYR1* reduced transpiration by decreasing the stomatal conductance without influencing photosynthesis under normal conditions. 35S:*PeCHYR1* poplar achieved higher photosynthesis and biomass accumulation along with reduced levels of MDA and REC under drought stress. Therefore, the *PeCHYR1* gene family will be conducive to further exploration of the mechanisms of drought tolerance, and we obtained a novel line of poplars with increased drought tolerance.

Experimental procedures

Plant materials

One-year-old seedlings of *P. euphratica* with heights of 15 cm were obtained from Yuli County, Korla, Xinjiang province, China (41° 19' 37.06" N, 86° 15' 44.78" E) in March, which lies in a warm temperate zone and has a continental desert climate. The average annual rainfall is 43 mm, and the average annual evaporation is 2700 mm. One-year-old seedlings of *P. euphratica* were transplanted in

This article is protected by copyright. All rights reserved.

individual pots (10 L) containing sandy soil (approximately 70% sand) and placed in a seed plot [light cycle: 16.0 h of light (06:00am–10:00pm); temperature (20 °C–24 °C)] at Hai Dian, Beijing, China (40°00'N, 116°20'E; 49 m above sea level). Potted *P. euphratica* were watered on the basis of evaporation demand and irrigated with 1 L of Hoagland nutrient solution every 2 weeks for 2 months before treatment (Li et al., 2011; Wang et al., 2016).

To analyze gene expression, similarly grown seedlings of *P. euphratica* (40–50 cm high, with 30–40 leaves) were subjected to ABA treatment and dehydration. The seedlings were treated with a solution containing 200.0 μM ABA (Sigma, A1049, USA) and 1 g TWEEN® 40 (Sigma, P1504-500 ML), which was sprayed on the leaves of *P. euphratica*. Meanwhile, the control was treated with 1 g TWEEN® 40 in ultrapure water. Dehydration treatment was applied to seedlings that had been removed from the soil and exposed to air with 70% RH at 23 °C under dim light for 12 h (Ma et al., 2010; Wang et al., 2016). To obtain complete plants, we gently pulled out each seedling and carefully washed the roots to remove the soil. For every test, leaves were separated from plants at the given time periods and promptly immersed in liquid nitrogen. We collected different organs and tissues of 2-month-old *P. euphratica*, including YL, young leaf; AL, adult leaf; OL, old leaf; Pe, petiole; X, xylem; and R, root at the same time and immediately immersed them in liquid nitrogen.

cDNA cloning of *PeCHYRI* from *P. euphratica*

Total RNA was extracted from the collected materials using the cetyltrimethylammonium bromide (CTAB) method (Chang et al., 1993). In the last step, potentially contaminating DNA was removed by treatment with DNase I. We used a NanoDrop 2000 Spectrophotometer (Thermo, West Palm

Beach, FL) to measure the quality and quantity of RNA. Two micrograms of total RNA was used for the reverse transcription reaction with a Tiangen FastQuant RT Kit (with gDNase) (Qiagen, Düsseldorf, Germany) according to the protocol. The cDNA of *PeCHYR1* was cloned by PCR, and the primers are displayed in supplement Table S1.

Quantitative real-time polymerase chain reaction (RT-qPCR) analysis

Twenty microliters of cDNA was diluted 1:10 with nuclease-free water. Each reaction contained 10 μ l SuperReal PreMix Plus (Tiangen Bio Inc., Beijing, China), 2 μ l ROX Reference Dye (Qiagen), 1 μ l cDNA [single-stranded circular DNA (sscDNA), corresponding to 10 ng of total RNA], 5.8 μ l nuclease-free water and 0.6 μ M of each primer. The cycling parameters were 95 °C for 15 min and then 45 cycles of 20 s at 95 °C and 60 s at 60 °C (Wang et al., 2014). RT-qPCR was performed with the ABI StepOnePlus Real-Time PCR System (ABI, Foster City, CA) according to the manufacturer's specifications. Each experiment was performed in 20 replicates (five biological replicates \times four technical replicates), and all primers used are displayed in Table S1. We used the software tool Primer Premier 6 to design primers. We calculated the relative expression level of *PeCHYR1* by the ratio = $(Et)_{\Delta CT} / (Er)_{\Delta CT}$ method (Wang et al., 2016; Wang et al., 2014).

Phylogenetic and domain analysis of *PeCHYR1*

Homologous amino acid sequences were obtained from the Phytozome database (<http://www.phytozome.net/>), and a phylogenetic tree of *PeCHYR1* was estimated with MEGA 7.

This article is protected by copyright. All rights reserved.

DNAMAN was used to examine the amino acid sequences and confirm the conserved structures. The sequence data from this study can be found in the Phytozome database (<http://www.phytozome.net/>). The accession numbers of the genes used are displayed in supplement Table S2.

Plasmid construction and genetic transformation of poplar

The *PeCHYR1* sequence was inserted into the *Sma*I and *Sac*I sites in the pCAMBIA-1301 vector. The construct was transformed into *Agrobacterium tumefaciens* (*EHA105*) and then transformed into poplar 84K using the leaf disc method (Hsu et al., 2011; Yao et al., 2016). The leaves of poplar 84K were incubated on substrate (pH 5.80) containing 0.020 mg/L thidiazuron (TDZ), 0.50 mg/L 6-benzylaminopurine (6-BA), 0.050 mg/L 1-naphthaleneacetic acid (NAA), 250 mg/L cefotaxime, 400 mg/L hygromycin and 0.80% (w/v) agar for shoot induction and selection. The regenerated shoots were individually detached from the callus and inserted into a rooting culture [1/2 MS substrate including 0.050 mg/L NAA, 250 mg/L cefotaxime, 150 mg/L hygromycin phosphotransferase and 0.9% (w/v) agar]. One-month-old regenerated seedlings were transplanted to small pots (12 cm length × 12 cm width × 12 cm height) with the same soil potting soil, turfy soil, and vermiculite 1:1:1) and then placed in a greenhouse [light cycle: 16.0 h of light, 8.0 h of dark; temperature: 24 °C (light)/20 °C (dark); relative humidity: 40%–45%].

Subcellular localization

To determine the subcellular localization of *PeCHYR1*, 35S:PeCHYR1-GFP fusion protein and 35S:HDEL-RFP fusion protein were transiently transfected together into tobacco leaves by the previously published protocol (Cui et al., 2012). Co-transformation of 35S:HDEL-RFP in combination with 35S:PeCHYR1-GFP was also performed in *Arabidopsis* leaf protoplasts by means of polyethylene glycol (PEG) treatment (Yoo et al., 2007). HDEL-RFP was used as an ER localization marker (Cui et al., 2012; Tian et al., 2013). The nuclear dye DAPI (10 mg/mL; Sigma-Aldrich) was applied to mark the nucleus. The transformed protoplasts and tobacco leaves were observed by laser confocal fluorescence microscopy (Leica TCS SP8). The LAS-AF software was used to record the images. The following lines of the argon ion laser were used: 488 nm for GFP, 584 nm for RFP, 380 nm for DAPI and 488 nm for chlorophyll. Fluorescence was detected at 495–515 nm for GFP, 575–590 nm for RFP, 430–450 nm for DAPI and 650 nm for chlorophyll.

Molecular verification of transgenic plants

Total DNA was extracted from every line using the CTAB method (Jiao et al., 2015; Verbylaite et al., 2010). We identified transformation lines by PCR using a forward primer (CAMV 35S promoter) and reverse primer (*PeCHYR1*). RT-qPCR was used for the analyses of *PeCHYR1* expression levels in different transgenic lines. Total RNA was obtained through the CTAB method from the leaves of WT and transgenic plants. Two micrograms of total RNA was used for the reverse transcription reaction with the Tiangen FastQuant RT Kit (with gDNase) (Qiagen, Düsseldorf, Germany), according to the

This article is protected by copyright. All rights reserved.

Accepted Article

manufacturer's instructions. qRT-PCR was performed with the ABI StepOnePlus Real-Time PCR System (ABI, Foster City, CA) according to the manufacturer's specifications. The internal control was a combination of two reference genes, *PeActin* and *PeUBQ*, which were screened previously (Bustin et al., 2009; Czechowski et al., 2005; Rao et al., 2013; Wang et al., 2014). The primers used are shown in Table S1.

Histochemical staining of GUS activity

Histochemical staining of GUS was handled as described previously (Lee et al., 2010). Briefly, isolated leaves were immersed at 37.0 °C in a mix including 2 mM 5-bromo-4-chloro-3-indolyl- β -D-glucuronide, 0.1 M sodium phosphate buffer (pH 7.0), 0.50 mM each of potassium ferri- and ferrocyanide, 10.0 mM EDTA (pH 7.0), and 0.10% Triton X-100 for 12 h. Then, the isolated leaves were incubated in 70% ethanol for 12 hours and photographed.

DAB staining

DAB staining was applied to detect the production of H₂O₂, as described previously (Ding et al., 2015a). The leaves of 2-month-old transgenic and WT plants were sprayed with 100 μ M ABA (Sigma, A1049, USA) and 1 g TWEEN® 40 (Sigma, P1504-500 ML) for 0, 1, 2, and 3 h and infiltrated in prepared 0.1 mg/mL DAB solution (Beyotime, ST033) in the dark for 8 h at 28 °C. Meanwhile, the controls were treated with 1 g TWEEN® 40 in ultrapure water. To remove

chlorophyll after DAB staining, the stained leaves were incubated in a solution of 70% ethanol, and then, they were photographed.

H₂O₂ detection in guard cells

A unique fluorescent dye, H₂DCF-DA (Sigma, D6883 HZB1212, USA), was used for H₂O₂ detection in the guard cells of the WT and transgenic plants (Huang et al., 2009). To observe transient ABA responses, the leaves of two-month-old seedlings (WT, *OXPeCHYRI*) were immersed in 50 mM H₂DCF-DA for 12 min at 25 °C in the dark. Then, the leaves were steeped in liquid 1/2 MS and treated with 100 μM ABA. Confocal measurements were taken every 30 s for 5 min of ABA stress in xyt mode. The amount of H₂DCF-DA fluorescence in guard cells was measured with Image-Pro Plus6 (Shi et al., 2013a).

Stomatal movement analysis

Stomatal movement tests were performed as described previously (Cui et al., 2015; Hsu et al., 2014b; Li et al., 2017), with slight amendments. Leaves from two-month-old seedlings (WT and *OXPeCHYRI*) were detached into wire netting and immediately fixed in liquid nitrogen. Then, the fixed samples were immediately put into an -80 °C Ultra-low Freeze Dryer (Biosafar-18A, Jiangsu, China (Mainland)) and fully dried for 24 hours. The fully dried sections were coated with gold using an Ion Sputter Coater (Eiko Engineering, IB-2, Japan). All observations were collected using a scanning electron microscope (Hitachi S-4700, Japan). For the ABA-induced stomatal closure experiments, the same parts of the leaves were immersed in stomata-opening solution (OS) containing 0.01 M KCl, 0.1 M CaCl₂, and 0.01 mM MES-KOH for 0.5 h in the dark and 2.0 h in the light, and

This article is protected by copyright. All rights reserved.

then 5 μM ABA was added. Then, stomata were fixed after 0, 1, and 2 h of ABA treatment. The stomatal images were collected using a Hitachi S-4700 scanning electron microscope. More than 100 guard cells from each sample were measured to derive the stomatal aperture.

Drought tolerance experiment

Fifteen 35S:*PeCHYR1* and five WT poplar plants grown in a greenhouse [light cycle: 16.0 h of light (06:00am–10:00pm); temperature (20 °C–24 °C)] were subjected to a short-term drought experiment. All of the plants (height 18–20 cm) were grown in suitably sized pots (12 cm \times 12 cm \times 12 cm), and each pot had a tray. All of the plants encountered drought treatment, in which the soil RWC was reduced from 70%. The control plants were kept in the same conditions, except that the soil RWC was maintained at 70%. A soil RWC of 70% induces no stress and is the most suitable environment for plant growth (Wang et al., 2016).

Physiological and biochemical analysis

A Li-Cor portable photosynthesis meter (LI-COR 6400) was applied to detect net photosynthetic rate, transpiration, and stomatal conductance in the ninth to eleventh leaves of 35S:*PeCHYR1* and WT poplar plants. We used the LI-COR 6400 portable photosynthesis analysis system to measure light curves under normal conditions of poplars grown in a greenhouse for 1.5 months. Twenty plants were measured. The experiment covered net CO₂ assimilation (A), Gs, transpiration, VPD and instantaneous WUE (A/transpiration). Photosynthetic light response curves were determined at

Accepted Article

photosynthetically active radiation (PAR) levels of 1500, 1200, 1000, 800, 600, 400, 200, 150, 100, 80, 50, 20, and 0 $\mu\text{m}^2/\text{s}$ with 450 $\mu\text{M}/\text{mol}$ external CO_2 , which was controlled by the LI-COR 6400 (Wang et al., 2016).

Measurement of H_2O_2 content and antioxidant enzyme activities

On the basis of the continuous changes of photosynthesis under drought stress, H_2O_2 and antioxidant enzyme activities were measured at 0, 3, and 7 days. Samples of 0.5 g fresh leaves were detached from the drought treatment and control plants at 0, 3, and 7 d and ground into a fine powder in liquid nitrogen. The plant extracts were isolated in 50 mM sodium phosphate buffer (pH 7.8) mixed thoroughly with 1 ml of 0.1% (w/v) titanium sulfate (in 20%, v/v H_2SO_4) for 10 min (Shi et al., 2012; Shi et al., 2014b). H_2O_2 content and the activities of antioxidant enzymes (SOD and POD) were measured using previously published protocols (Shi et al., 2014a). The levels of H_2O_2 , SOD and POD were expressed as μM ($\text{g} \cdot \text{fresh weight (FW)}^{-1}$). These experiments consisted of 20 replicates (4 biological replicates \times 5 technical replicates) under identical conditions.

MDA level and relative electrical conductance (REC) assay

Materials detached from drought-stressed plants at 0 and 7d were measured to determine MDA level and REC using previously published protocols (Shi et al., 2013b; Shi et al., 2014b). MDA (mol/g FW) = $(6.45 \cdot (A_{532} - A_{600}) - 0.56 \cdot A_{450}) \cdot 9 \cdot V_t / V_s - 1 \text{FW}$ [Vt: total volume of extract (ml); Vs: extract volume was determined with (ml)]. The electrical conductivity of the supernatant (L1) was detected using a DDS-307 Conductivity Meter (Leici-DDS-307A, Shanghai, China). These experiments were independently replicated 6 times under identical conditions.

This article is protected by copyright. All rights reserved.

Analysis of chlorophyll fluorescence and content

Photosynthetic activity in the 10th–13th leaves of transgenic and WT plants was monitored by photosynthetic efficiency of photosystem II (Fv/Fm) values, which reflect the potential maximum light energy conversion efficiency of plants, using a PAM chlorophyll fluorometer (PAM100) after 20 min of dark adaptation. Chlorophyll content was detected in the 10th–13th leaves of transgenic and WT plants using a portable chlorophyll meter (SPAD-502Plus, Konica Minolta, Japan). Total chlorophyll contents were measured at 70% soil RWC and 45% soil RWC.

Long-term drought experiment

Drought stress was implemented in transgenic and WT poplar plants by maintaining the soil RWC at 45% for 32 days. Control plants were maintained at a soil RWC of 70%. A soil RWC of 70% causes no stress and is a suitable environment for plant growth (Wang et al., 2016). The transgenic and WT poplar plants (five poplars per line under various treatment) were grown in a greenhouse [light cycle: 16.0 h of light (06:00am–10:00pm); temperature (20 °C–24 °C)], all plants (height 40–43 cm) were kept in suitably sized pots (volume, 15 L), and every pot had a tray. The containers were weighed daily, and lost water was supplemented. We measured the height, shoot biomass, and root biomass of each poplar every 7 days. After 32 days, chlorophyll content, leaf RWC, and maximum photosynthetic efficiency of photosystem II (Fv/Fm) were measured.

Relative water content (RWC)

The 10th–13th leaves were detached and used for RWC measurements. In short, we determined the leaf FW (removed fresh leaves were weighed), leaf turgid weight (TW, leaves were measured after submerging in water for 8 h), and leaf dry weight (DW, leaves were measured after drying at 80 °C for 72 h). The RWC was calculated as $(FW - DW) / (TW - DW) * 100\%$.

Statistical analysis

The experimental data were subjected to analysis of variance by Statistical Product and Service Solutions 17.0 (SPSS). For statistical analyses, Student's t-test was used to generate every P value (one-way analysis of variance: $P < 0.05$, *; $P < 0.01$, **). Both one-way and two-way analyses of variance were used to determine significance. The data were normalized, and all samples were normally distributed with homogeneity of variance.

Acknowledgements

This study was supported by grants from the Ministry of Science and Technology of China (2015BAD07B01, 2013AA102701) and the National Natural Science Foundation of China (31270656, 31570308). The authors declare no conflict of interest.

References

- Ahmad, P., Sarwat, M. and Sharma, S. (2008) Reactive oxygen species, antioxidants and signaling in plants. *J Plant Biol* **51**, 167-173.
- Bae, H., Kim, S.K., Cho, S.K., Kang, B.G. and Kim, W.T. (2011) Overexpression of OsRDCP1, a rice RING domain-containing E3 ubiquitin ligase, increased tolerance to drought stress in rice (*Oryza sativa* L.). *Plant Science* **180**, 775-782.
- Bhattacharjee, S. (2012) The Language of Reactive Oxygen Species Signaling in Plants. *Journal of Botany* **2012**, 1-22.
- Bright, J., Desikan, R., Hancock, J.T., Weir, I.S. and Neill, S.J. (2006) ABA-induced NO generation and stomatal closure in *Arabidopsis* are dependent on H₂O₂ synthesis. *Plant Journal* **45**, 113-122.
- Bussis, D., von Groll, U., Fisahn, J. and Altmann, T. (2006) Stomatal aperture can compensate altered stomatal density in *Arabidopsis thaliana* at growth light conditions. *Functional Plant Biology* **33**, 1037-1043.
- Bustin, S.A., Benes, V., Garson, J.A., Hellems, J., Huggett, J., Kubista, M., Mueller, R., Nolan, T., Pfaffl, M.W., Shipley, G.L., Vandesompele, J. and Wittwer, C.T. (2009) The MIQE Guidelines: Minimum Information for Publication of Quantitative Real-Time PCR Experiments. *Clin Chem* **55**, 611-622.
- Cayrol, C., Lacroix, C., Mathe, C., Ecochard, V., Ceribelli, M., Loreau, E., Lazar, V., Dessen, P., Mantovani, R., Aguilar, L. and Girard, J.P. (2007) The THAP-zinc finger protein THAP1 regulates endothelial cell proliferation through modulation of pRB/E2F cell-cycle target genes. *Blood* **109**, 584-594.

Chaerle, L., Saibo, N. and Van Der Straeten, D. (2005) Tuning the pores: towards engineering plants for improved water use efficiency. *Trends in Biotechnology* **23**, 308-315.

Chang, S., Puryear, J. and Cairney, J. (1993) A simple and efficient method for isolating RNA from pine trees. *Plant Mol Biol Rep* **11**, 113-116.

Chen, J.H., Xue, B., Xia, X.L. and Yin, W.L. (2013) A novel calcium-dependent protein kinase gene from *Populus euphratica*, confers both drought and cold stress tolerance. *Biochem Bioph Res Co* **441**, 630-636.

Cui, F., Liu, L.J., Zhao, Q.Z., Zhang, Z.H., Li, Q.L., Lin, B.Y., Wu, Y.R., Tang, S.Y. and Xie, Q. (2012) Arabidopsis Ubiquitin Conjugase UBC32 Is an ERAD Component That Functions in Brassinosteroid-Mediated Salt Stress Tolerance. *Plant Cell* **24**, 233-244.

Cui, L.G., Shan, J.X., Shi, M., Gao, J.P. and Lin, H.X. (2015) DCA1 Acts as a Transcriptional Co-activator of DST and Contributes to Drought and Salt Tolerance in Rice. *Plos Genet* **11**.

Czechowski, T., Stitt, M., Altmann, T., Udvardi, M.K. and Scheible, W.R. (2005) Genome-wide identification and testing of superior reference genes for transcript normalization in Arabidopsis. *Plant Physiology* **139**, 5-17.

Danquah, A., de Zelicourt, A., Colcombet, J. and Hirt, H. (2014) The role of ABA and MAPK signaling pathways in plant abiotic stress responses. *Biotechnol Adv* **32**, 40-52.

Dash, M., Yordanov, Y.S., Georgieva, T., Tschaplinski, T.J., Yordanova, E. and Busov, V. (2017) Poplar PtabZIP1-like enhances lateral root formation and biomass growth under drought stress. *Plant J* **89**, 692-705.

de la Garma, J.G., Fernandez-Garcia, N., Bardisi, E., Pallol, B., Rubio-Asensio, J.S., Bru, R. and Olmos, E. (2015)

New insights into plant salt acclimation: the roles of vesicle trafficking and reactive oxygen species signalling in mitochondria and the endomembrane system. *New Phytol* **205**, 216-239.

Dietz, K.J., Mittler, R. and Noctor, G. (2016) Recent Progress in Understanding the Role of Reactive Oxygen Species in Plant Cell Signaling. *Plant Physiology* **171**, 1535-1539.

Ding, M.Q., Hou, P.C., Shen, X., Wang, M.J., Deng, S.R., Sun, J., Xiao, F., Wang, R.G., Zhou, X.Y., Lu, C.F., Zhang, D.Q., Zheng, X.J., Hu, Z.M. and Chen, S.L. (2010) Salt-induced expression of genes related to Na⁺/K⁺ and ROS homeostasis in leaves of salt-resistant and salt-sensitive poplar species. *Plant Mol Biol* **73**, 251-269.

Ding, S., Zhang, B. and Qin, F. (2015a) Arabidopsis RZFP34/CHYR1, a Ubiquitin E3 Ligase, Regulates Stomatal Movement and Drought Tolerance via SnRK2.6-Mediated Phosphorylation. *Plant Cell* **27**, 3228-3244.

Ding, S.C., Zhang, B. and Qin, F. (2015b) Arabidopsis RZFP34/CHYR1, a Ubiquitin E3 Ligase, Regulates Stomatal Movement and Drought Tolerance via SnRK2.6-Mediated Phosphorylation. *Plant Cell* **27**, 3228-3244.

Esposito, S., Baglivo, I., Malgieri, G., Russo, L., Zaccaro, L., D'Andrea, L.D., Mammucari, M., Di Blasio, B., Isernia, C., Fattorusso, R. and Pedone, P.V. (2006) A novel type of zinc finger DNA binding domain in the *Agrobacterium tumefaciens* transcriptional regulator Ros. *Biochemistry* **45**, 10394-10405.

Feng, F., Ding, F. and Tyree, M.T. (2015) Investigations Concerning Cavitation and Frost Fatigue in Clonal 84K Poplar Using High-Resolution Cavitron Measurements. *Plant Physiology* **168**, 144-U850.

- Gudesblat, G.E., Iusem, N.D. and Morris, P.C. (2007) Guard cell-specific inhibition of Arabidopsis MPK3 expression causes abnormal stomatal responses to abscisic acid and hydrogen peroxide. *New Phytologist* **173**, 713-721.
- Hetherington, A.M. and Woodward, F.I. (2003) The role of stomata in sensing and driving environmental change. *Nature* **424**, 901-908.
- Hsu, C.Y., Adams, J.P., Kim, H.J., No, K., Ma, C.P., Strauss, S.H., Drnevich, J., Vandervelde, L., Ellis, J.D., Rice, B.M., Wickett, N., Gunter, L.E., Tuskan, G.A., Brunner, A.M., Page, G.P., Barakat, A., Carlson, J.E., dePamphilis, C.W., Luthe, D.S. and Yuceer, C. (2011) FLOWERING LOCUS T duplication coordinates reproductive and vegetative growth in perennial poplar. *P Natl Acad Sci USA* **108**, 10756-10761.
- Hsu, K.H., Liu, C.C., Wu, S.J., Kuo, Y.Y., Lu, C.A., Wu, C.R., Lian, P.J., Hong, C.Y., Ke, Y.T., Huang, J.H. and Yeh, C.H. (2014a) Expression of a gene encoding a rice RING zinc-finger protein, OsRZFP34, enhances stomata opening. *Plant Mol Biol* **86**, 125-137.
- Hsu, K.H., Liu, C.C., Wu, S.J., Kuo, Y.Y., Lu, C.A., Wu, C.R., Lian, P.J., Hong, C.Y., Ke, Y.T., Huang, J.H. and Yeh, C.H. (2014b) Expression of a gene encoding a rice RING zinc-finger protein, OsRZFP34, enhances stomata opening. *Plant Mol Biol* **86**, 125-137.
- Hua, D.P., Wang, C., He, J.N., Liao, H., Duan, Y., Zhu, Z.Q., Guo, Y., Chen, Z.Z. and Gong, Z.Z. (2012) A Plasma Membrane Receptor Kinase, GHR1, Mediates Abscisic Acid- and Hydrogen Peroxide-Regulated Stomatal Movement in Arabidopsis. *Plant Cell* **24**, 2546-2561.

- Huang, S.B., Van Aken, O., Schwarzlander, M., Belt, K. and Millar, A.H. (2016) The Roles of Mitochondrial Reactive Oxygen Species in Cellular Signaling and Stress Response in Plants. *Plant Physiology* **171**, 1551-1559.
- Huang, X.Y., Chao, D.Y., Gao, J.P., Zhu, M.Z., Shi, M. and Lin, H.X. (2009) A previously unknown zinc finger protein, DST, regulates drought and salt tolerance in rice via stomatal aperture control. *Genes Dev* **23**, 1805-1817.
- Jiang, M. and Zhang, J. (2002) Water stress-induced abscisic acid accumulation triggers the increased generation of reactive oxygen species and up-regulates the activities of antioxidant enzymes in maize leaves. *J Exp Bot* **53**, 2401-2410.
- Jiao, L.C., Liu, X.L., Jiang, X.N. and Yin, Y.F. (2015) Extraction and amplification of DNA from aged and archaeological *Populus euphratica* wood for species identification. *Holzforschung* **69**, 925-931.
- Jin, Y.L., Tang, R.J., Wang, H.H., Jiang, C.M., Bao, Y., Yang, Y., Liang, M.X., Sun, Z.C., Kong, F.J., Li, B. and Zhang, H.X. (2017) Overexpression of *Populus trichocarpa* CYP85A3 promotes growth and biomass production in transgenic trees. *Plant Biotechnol J* **15**, 1309-1321.
- Joshi-Saha, A., Valon, C. and Leung, J. (2011) Abscisic Acid Signal off the STARTing Block. *Mol Plant* **4**, 562-580.
- Karuppanapandian, T., Moon, J.C., Kim, C., Manoharan, K. and Kim, W. (2011) Reactive oxygen species in plants: their generation, signal transduction, and scavenging mechanisms. *Aust J Crop Sci* **5**, 709-725.

Ke, Q.B., Wang, Z., Ji, C.Y., Jeong, J.C., Lee, H.S., Li, H.B., Xu, B.C., Deng, X.P. and Kwak, S.S. (2016)

Transgenic poplar expressing *codA* exhibits enhanced growth and abiotic stress tolerance. *Plant Physiol Bioch* **100**, 75-84.

Kitajima, M. and Butler, W.L. (1975) Quenching of chlorophyll fluorescence and primary photochemistry in chloroplasts by dibromothymoquinone. *Biochim Biophys Acta* **376**, 105-115.

Kobayashi, Y., Murata, M., Minami, H., Yamamoto, S., Kagaya, Y., Hobo, T., Yamamoto, A. and Hattori, T. (2005) Abscisic acid-activated SNRK2 protein kinases function in the gene-regulation pathway of ABA signal transduction by phosphorylating ABA response element-binding factors. *Plant J* **44**, 939-949.

Kohler, B., Hills, A. and Blatt, M.R. (2003) Control of guard cell ion channels by hydrogen peroxide and abscisic acid indicates their action through alternate signaling pathways. *Plant Physiology* **131**, 385-388.

Kolla, V.A., Vavasseur, A. and Raghavendra, A.S. (2007) Hydrogen peroxide production is an early event during bicarbonate induced stomatal closure in abaxial epidermis of Arabidopsis. *Planta* **225**, 1421-1429.

Kreslavski, V.D., Los, D.A., Allakhverdiev, S.I. and Kuznetsov, V.V. (2012) Signaling Role of Reactive Oxygen Species in Plants under Stress. *Russ J Plant Physiol+* **59**, 141-154.

Kuge, S., Tachibana, T., Iwai, K. and Naganuma, A. (2010) Peroxiredoxin-induced sensing and transduction of redox signal in response to oxidative stress and metabolism in yeast cells. *Journal of Pharmacological Sciences* **112**, 22p-22p.

Larcheveque, M., Maurel, M., Desrochers, A. and Larocque, G.R. (2011) How does drought tolerance compare between two improved hybrids of balsam poplar and an unimproved native species? *Tree Physiol* **31**, 240-249.

This article is protected by copyright. All rights reserved.

Lee, J.H. and Kim, W.T. (2011) Regulation of abiotic stress signal transduction by E3 ubiquitin ligases in

Arabidopsis. *Molecules and Cells* **31**, 201-208.

Lee, O.R., Kim, S.J., Kim, H.J., Hong, J.K., Ryu, S.B., Lee, S.H., Ganguly, A. and Cho, H.T. (2010)

Phospholipase A(2) Is Required for PIN-FORMED Protein Trafficking to the Plasma Membrane in the
Arabidopsis Root. *Plant Cell* **22**, 1812-1825.

Leng, R.P., Lin, Y.P., Ma, W.L., Wu, H., Lemmers, B., Chung, S., Parant, J.M., Lozano, G., Hakem, R. and

Benchimol, S. (2003) Pirh2, a p53-induced ubiquitin-protein ligase, promotes p53 degradation. *Cell* **112**,
779-791.

Li, B.S., Qin, Y.R., Duan, H., Yin, W.L. and Xia, X.L. (2011) Genome-wide characterization of new and drought

stress responsive microRNAs in *Populus euphratica*. *J Exp Bot* **62**, 3765-3779.

Li, J.J., Li, Y., Yin, Z.G., Jiang, J.H., Zhang, M.H., Guo, X., Ye, Z.J., Zhao, Y., Xiong, H.Y., Zhang, Z.Y., Shao,

Y.J., Jiang, C.H., Zhang, H.L., An, G.H., Paek, N.C., Ali, J. and Li, Z.C. (2017) OsASR5 enhances
drought tolerance through a stomatal closure pathway associated with ABA and H₂O₂ signalling in rice.
Plant Biotechnol J **15**, 183-196.

Lim, C.W., Baek, W. and Lee, S.C. (2017) The Pepper RING-Type E3 Ligase CaAIRF1 Regulates ABA and

Drought Signaling via CaADIP1 Protein Phosphatase Degradation. *Plant Physiology* **173**, 2323-2339.

Lim, S.D., Cho, H.Y., Park, Y.C., Ham, D.J., Lee, J.K. and Jang, C.S. (2013a) The rice RING finger E3 ligase,

OshCII, drives nuclear export of multiple substrate proteins and its heterogeneous overexpression
enhances acquired thermotolerance. *Journal of Experimental Botany* **64**, 2899-2914.

Lim, S.D., Hwang, J.G., Jung, C.G., Hwang, S.G., Moon, J.C. and Jang, C.S. (2013b) Comprehensive Analysis of the Rice RING E3 Ligase Family Reveals Their Functional Diversity in Response to Abiotic stress. *DNA Research* **20**, 299-314.

Liu, J., Zhang, C., Wei, C., Liu, X., Wang, M., Yu, F., Xie, Q. and Tu, J. (2016) The RING Finger Ubiquitin E3 Ligase OsHTAS Enhances Heat Tolerance by Promoting H₂O₂-Induced Stomatal Closure in Rice. *Plant Physiol* **170**, 429-443.

Lyzenga, W.J. and Stone, S.L. (2012) Abiotic stress tolerance mediated by protein ubiquitination. *Journal of Experimental Botany* **63**, 599-616.

Ma, H.S., Liang, D., Shuai, P., Xia, X.L. and Yin, W.L. (2010) The salt- and drought-inducible poplar GRAS protein SCL7 confers salt and drought tolerance in Arabidopsis thaliana. *J Exp Bot* **61**, 4011-4019.

Ma, T., Wang, J.Y., Zhou, G.K., Yue, Z., Hu, Q.J., Chen, Y., Liu, B.B., Qiu, Q., Wang, Z., Zhang, J., Wang, K., Jiang, D.C., Gou, C.Y., Yu, L.L., Zhan, D.L., Zhou, R., Luo, W.C., Ma, H., Yang, Y.Z., Pan, S.K., Fang, D.M., Luo, Y.D., Wang, X., Wang, G.N., Wang, J., Wang, Q., Lu, X., Chen, Z., Liu, J.C., Lu, Y., Yin, Y., Yang, H.M., Abbott, R.J., Wu, Y.X., Wan, D.S., Li, J., Yin, T.M., Lascoux, M., DiFazio, S.P., Tuskan, G.A., Wang, J. and Liu, J.Q. (2013) Genomic insights into salt adaptation in a desert poplar. *Nat Commun* **4**.

Ma, X.F., Wang, W.M., Bittner, F., Schmidt, N., Berkey, R., Zhang, L.L., King, H., Zhang, Y., Feng, J.Y., Wen, Y.Q., Tan, L.Q., Li, Y., Zhang, Q., Deng, Z.N., Xiong, X.Y. and Xiao, S.Y. (2016) Dual and Opposing Roles of Xanthine Dehydrogenase in Defense-Associated Reactive Oxygen Species Metabolism in Arabidopsis. *Plant Cell* **28**, 1108-1126.

McAinsh, M.R., Clayton, H., Mansfield, T.A. and Hetherington, A.M. (1996) Changes in Stomatal Behavior and Guard Cell Cytosolic Free Calcium in Response to Oxidative Stress. *Plant Physiol* **111**, 1031-1042.

McDowall (2007) <Zinc finger.pdf>. <http://www.ebi.ac.uk>.

Moradi, P. (2016) Key plant products and common mechanisms utilized by plants in water deficit stress responses. *Bot Sci* **94**, 657-671.

Niinemets, U. (2002) Stomatal conductance alone does not explain the decline in foliar photosynthetic rates with increasing tree age and size in *Picea abies* and *Pinus sylvestris*. *Tree Physiol* **22**, 515-535.

Ning, Y.S., Wang, R.Y., Shi, X.T., Zhou, X.P. and Wang, G.L. (2016) A Layered Defense Strategy Mediated by Rice E3 Ubiquitin Ligases against Diverse Pathogens. *Mol Plant* **9**, 1096-1098.

Park, C., Lim, C.W., Baek, W. and Lee, S.C. (2015) RING Type E3 Ligase CaAIR1 in Pepper Acts in the Regulation of ABA Signaling and Drought Stress Response. *Plant Cell Physiol* **56**, 1808-1819.

Pearcy, R.W. and Seemann, J.R. (1990) Photosynthetic Induction State of Leaves in a Soybean Canopy in Relation to Light Regulation of Ribulose-1-5-Bisphosphate Carboxylase and Stomatal Conductance. *Plant Physiology* **94**, 628-633.

Pei, Z.M., Murata, Y., Benning, G., Thomine, S., Klusener, B., Allen, G.J., Grill, E. and Schroeder, J.I. (2000) Calcium channels activated by hydrogen peroxide mediate abscisic acid signalling in guard cells. *Nature* **406**, 731-734.

Pitzschke, A., Forzani, C. and Hirt, H. (2006) Reactive oxygen species signaling in plants. *Antioxid Redox Sign* **8**, 1757-1764.

Accepted Article

Qin, L., Kang, W.-h., Qi, Y.-l., Zhang, Z.-w. and Wang, N. (2016) The influence of silicon application on growth and photosynthesis response of salt stressed grapevines (*Vitis vinifera* L.). *Acta Physiologiae Plantarum* **38**, 1-9.

Rao, X.Y., Lai, D.J. and Huang, X.L. (2013) A New Method for Quantitative Real-Time Polymerase Chain Reaction Data Analysis. *J Comput Biol* **20**, 703-711.

Shi, H., Ye, T., Chen, F., Cheng, Z., Wang, Y., Yang, P., Zhang, Y. and Chan, Z. (2013a) Manipulation of arginase expression modulates abiotic stress tolerance in Arabidopsis: effect on arginine metabolism and ROS accumulation. *J Exp Bot* **64**, 1367-1379.

Shi, H., Ye, T., Zhu, J.K. and Chan, Z. (2014a) Constitutive production of nitric oxide leads to enhanced drought stress resistance and extensive transcriptional reprogramming in Arabidopsis. *J Exp Bot* **65**, 4119-4131.

Shi, H.T., Wang, Y.P., Cheng, Z.M., Ye, T.T. and Chan, Z.L. (2012) Analysis of Natural Variation in Bermudagrass (*Cynodon dactylon*) Reveals Physiological Responses Underlying Drought Tolerance. *Plos One* **7**.

Shi, H.T., Ye, T.T., Chen, F.F., Cheng, Z.M., Wang, Y.P., Yang, P.F., Zhang, Y.S. and Chan, Z.L. (2013b) Manipulation of arginase expression modulates abiotic stress tolerance in Arabidopsis: effect on arginine metabolism and ROS accumulation. *J Exp Bot* **64**, 1367-1379.

Shi, H.T., Ye, T.T., Zhu, J.K. and Chan, Z.L. (2014b) Constitutive production of nitric oxide leads to enhanced drought stress resistance and extensive transcriptional reprogramming in Arabidopsis. *J Exp Bot* **65**, 4119-4131.

- Shinozaki, K. and Yamaguchi-Shinozaki, K. (2007) Gene networks involved in drought stress response and tolerance. *J Exp Bot* **58**, 221-227.
- Sirichandra, C., Gu, D., Hu, H.C., Davanture, M., Lee, S., Djaoui, M., Valot, B., Zivy, M., Leung, J., Merlot, S. and Kwak, J.M. (2009) Phosphorylation of the Arabidopsis AtrbohF NADPH oxidase by OST1 protein kinase. *FEBS Lett* **583**, 2982-2986.
- Sterky, F., Bhalerao, R.R., Unneberg, P., Segerman, B., Nilsson, P., Brunner, A.M., Charbonnel-Campaa, L., Lindvall, J.J., Tandré, K., Strauss, S.H., Sundberg, B., Gustafsson, P., Uhlen, M., Bhalerao, R.P., Nilsson, O., Sandberg, G., Karlsson, J., Lundeberg, J. and Jansson, S. (2004) A *Populus* EST resource for plant functional genomics. *Proceedings of the National Academy of Sciences of the United States of America* **101**, 13951-13956.
- Stone, S.L., Hauksdottir, H., Troy, A., Herschleb, J., Kraft, E. and Callis, J. (2005) Functional analysis of the RING-type ubiquitin ligase family of Arabidopsis. *Plant Physiology* **137**, 13-30.
- Tang, S., Liang, H.Y., Yan, D.H., Zhao, Y., Han, X., Carlson, J.E., Xia, X.L. and Yin, W.L. (2013) *Populus euphratica*: the transcriptomic response to drought stress. *Plant Mol Biol* **83**, 539-557.
- Tian, L.H., Dai, L.L., Yin, Z.J., Fukuda, M., Kumamaru, T., Dong, X.B., Xu, X.P. and Qu, L.Q. (2013) Small GTPase Sar1 is crucial for proglutelin and -globulin export from the endoplasmic reticulum in rice endosperm. *J Exp Bot* **64**, 2831-2845.
- Tuskan, G.A., DiFazio, S., Jansson, S., Bohlmann, J., Grigoriev, I., Hellsten, U., Putnam, N., Ralph, S., Rombauts, S., Salamov, A., Schein, J., Sterck, L., Aerts, A., Bhalerao, R.R., Bhalerao, R.P., Blaudez, D., Boerjan, W., Brun, A., Brunner, A., Busov, V., Campbell, M., Carlson, J., Chalot, M., Chapman, J.,

Chen, G.L., Cooper, D., Coutinho, P.M., Couturier, J., Covert, S., Cronk, Q., Cunningham, R., Davis, J., Degroeve, S., Dejardin, A., Depamphilis, C., Detter, J., Dirks, B., Dubchak, I., Duplessis, S., Ehlting, J., Ellis, B., Gendler, K., Goodstein, D., Gribskov, M., Grimwood, J., Groover, A., Gunter, L., Hamberger, B., Heinze, B., Helariutta, Y., Henrissat, B., Holligan, D., Holt, R., Huang, W., Islam-Faridi, N., Jones, S., Jones-Rhoades, M., Jorgensen, R., Joshi, C., Kangasjarvi, J., Karlsson, J., Kelleher, C., Kirkpatrick, R., Kirst, M., Kohler, A., Kalluri, U., Larimer, F., Leebens-Mack, J., Leple, J.C., Locascio, P., Lou, Y., Lucas, S., Martin, F., Montanini, B., Napoli, C., Nelson, D.R., Nelson, C., Nieminen, K., Nilsson, O., Pereda, V., Peter, G., Philippe, R., Pilate, G., Poliakov, A., Razumovskaya, J., Richardson, P., Rinaldi, C., Ritland, K., Rouze, P., Ryaboy, D., Schmutz, J., Schrader, J., Segerman, B., Shin, H., Siddiqui, A., Sterky, F., Terry, A., Tsai, C.J., Uberbacher, E., Unneberg, P., Vahala, J., Wall, K., Wessler, S., Yang, G., Yin, T., Douglas, C., Marra, M., Sandberg, G., Van de Peer, Y. and Rokhsar, D. (2006) The genome of black cottonwood, *Populus trichocarpa* (Torr. & Gray). *Science* **313**, 1596-1604.

Vahisalu, T., Kollist, H., Wang, Y.F., Nishimura, N., Chan, W.Y., Valerio, G., Lamminmaki, A., Brosche, M., Moldau, H., Desikan, R., Schroeder, J.I. and Kangasjarvi, J. (2008) SLAC1 is required for plant guard cell S-type anion channel function in stomatal signalling. *Nature* **452**, 487-U415.

Verbylaite, R., Beisy, P., Rimasa, V. and Kuusiene, S. (2010) Comparison of Ten DNA Extraction Protocols from Wood of European Aspen (*Populus tremula* L.). *Balt For* **16**, 35-42.

Wang, C., Liu, S., Dong, Y., Zhao, Y., Geng, A., Xia, X. and Yin, W. (2016) PdEPF1 regulates water-use efficiency and drought tolerance by modulating stomatal density in poplar. *Plant Biotechnol J* **14**, 849-860.

Accepted Article

Wang, H.L., Chen, J.H., Tian, Q.Q., Wang, S., Xia, X.L. and Yin, W.L. (2014) Identification and validation of reference genes for *Populus euphratica* gene expression analysis during abiotic stresses by quantitative real-time PCR. *Physiol Plantarum* **152**, 529-545.

Wang, R.G., Chen, S.L., Zhou, X.Y., Shen, X., Deng, L., Zhu, H.J., Shao, J., Shi, Y., Dai, S.X., Fritz, E., Huttermann, A. and Polle, A. (2008) Ionic homeostasis and reactive oxygen species control in leaves and xylem sap of two poplars subjected to NaCl stress. *Tree Physiology* **28**, 947-957.

Wege, S., De Angeli, A., Droillard, M.J., Kroniewicz, L., Merlot, S., Cornu, D., Gambale, F., Martinoia, E., Barbier-Brygoo, H., Thomine, S., Leonhardt, N. and Filleur, S. (2014) Phosphorylation of the vacuolar anion exchanger AtCLCa is required for the stomatal response to abscisic acid. *Sci Signal* **7**.

Willekens H, C.S., Davey M, Schraudner M, Langebartels C, Van Montagu M, Inzé D, Van Camp W (1997) <Willekens_et_al-1997-The_EMBO_Journal.pdf>. *EMBO J*.

Windt, C.W., Vergeldt, F.J., De Jager, P.A. and Van As, H. (2006) MRI of long-distance water transport: a comparison of the phloem and xylem flow characteristics and dynamics in poplar, castor bean, tomato and tobacco. *Plant Cell Environ* **29**, 1715-1729.

Winter, K. and Lesch, M. (1992) Diurnal changes in chlorophylla fluorescence and carotenoid composition in *Opuntia ficus-indica*, a CAM plant, and in three C3 species in Portugal during summer. *Oecologia* **91**, 505-510.

Wooff, J., Pastushok, L., Hanna, M., Fu, Y. and Xiao, W. (2004) The TRAF6 RING finger domain mediates physical interaction with Ubc13. *Febs Letters* **566**, 229-233.

Xing, Y., Jia, W. and Zhang, J. (2008) AtMKK1 mediates ABA-induced CAT1 expression and H₂O₂ production via AtMPK6-coupled signaling in Arabidopsis. *Plant J* **54**, 440-451.

Yan, D.H., Fenning, T., Tang, S., Xia, X.L. and Yin, W.L. (2012) Genome-wide transcriptional response of *Populus euphratica* to long-term drought stress. *Plant Sci* **195**, 24-35.

Yang, W., Liu, X.D., Chi, X.J., Wu, C.A., Li, Y.Z., Song, L.L., Liu, X.M., Wang, Y.F., Wang, F.W., Zhang, C.A., Liu, Y., Zong, J.M. and Li, H.Y. (2011) Dwarf apple MbdDREB1 enhances plant tolerance to low temperature, drought, and salt stress via both ABA-dependent and ABA-independent pathways. *Planta* **233**, 219-229.

Yao, W., Wang, S., Zhou, B. and Jiang, T. (2016) Transgenic poplar overexpressing the endogenous transcription factor ERF76 gene improves salinity tolerance. *Tree Physiol* **36**, 896-908.

Yin, C.Y., Duan, B.L., Wang, X. and Li, C.Y. (2004) Morphological and physiological responses of two contrasting Poplar species to drought stress and exogenous abscisic acid application. *Plant Science* **167**, 1091-1097.

Yoo, S.D., Cho, Y.H. and Sheen, J. (2007) Arabidopsis mesophyll protoplasts: a versatile cell system for transient gene expression analysis. *Nat Protoc* **2**, 1565-1572.

You, J., Zong, W., Li, X., Ning, J., Hu, H., Li, X., Xiao, J. and Xiong, L. (2013) The SNAC1-targeted gene OsSRO1c modulates stomatal closure and oxidative stress tolerance by regulating hydrogen peroxide in rice. *J Exp Bot* **64**, 569-583.

Yu, F.F., Wu, Y.R. and Xie, Q. (2016) Ubiquitin-Proteasome System in ABA Signaling: From Perception to Action. *Mol Plant* **9**, 21-33.

This article is protected by copyright. All rights reserved.

Zhang, X., Miao, Y.C., An, G.Y., Zhou, Y., Shangguan, Z.P., Gao, J.F. and Song, C.P. (2001) K⁺ channels inhibited by hydrogen peroxide mediate abscisic acid signaling in *Vicia* guard cells. *Cell Res* **11**, 195-202.

Zhao, Y., Sun, J., Xu, P., Zhang, R. and Li, L. (2014) Intron-mediated alternative splicing of WOOD-ASSOCIATED NAC TRANSCRIPTION FACTOR1B regulates cell wall thickening during fiber development in *Populus* species. *Plant Physiol* **164**, 765-776.

Zhu, J.K. (2016) Abiotic Stress Signaling and Responses in Plants. *Cell* **167**, 313-324.

Figure legends

Figure 1. Amino acid sequence alignment and phylogenetic tree of different CHYR1 protein family members. (a) Phylogenetic analysis of the *PeCHYR1* homologs from *Populus*, *Amborella*, *Eucalyptus*, *Malus*, *Prunus*, *Theobroma*, *Picea*, *Pinus*, *Salix*, *Arabidopsis*, rice, and maize. (b) Multiple alignment of the amino acid sequences of CHYR1 proteins from *Populus*, *Salix*, and *Arabidopsis*.

Figure 2. *PeCHYR1* expression patterns in different tissues and under different treatments. (a) Relative expression levels of the *PeCHYR1* gene in different tissues of *P. euphratica*. Young leaf; Adult leaf; Old leaf; Petiole; Xylem; Root. (b) Transcript levels of *PeCHYR1* were measured by RT-qPCR in response to dehydration. (c) Transcript levels of *PeCHYR1* were measured by RT-qPCR in response to ABA. (d) DAB staining in WT seedling leaves during ABA treatment. Error bars are means \pm SE (n = 20). Asterisks denote significant differences: *, P \leq 0.05; **, P \leq 0.01.

Figure 3. PeCHYR1 is targeted to the endoplasmic reticulum (ER) and nucleus.

(a) Subcellular localization of 35S: GFP and 35S:PeCHYR1-GFP in transiently expressed tobacco leaves. HDEL-RFP was used as an ER localization marker fused with red fluorescent protein (RFP). The nuclear dye DAPI (blue) was applied to mark the nucleus. Bar = 10 μ m. (b) Subcellular localization of 35S: GFP and 35S:PeCHYR1-GFP in transiently expressed Arabidopsis leaf protoplasts. 35S: HDE-RFP was co-transformed with 35S:PeCHYR1-GFP to verify the ER localization of PeCHYR1. The nuclear dye DAPI (blue) was applied to mark the nucleus. Bar = 5 μ m.

Figure 4. *PeCHYR1* promotes ABA-induced stomatal closure via ROS production. (a) DAB staining shows different levels of ABA-induced ROS production in the leaves of WT and 35S:*PeCHYR1*. (b) Representative confocal images of 200 μ M ABA-induced H₂O₂ production (10 min) in the guard cells of WT poplar and transgenic lines colored with H2DCFDA. (c) Quantification of H₂O₂ production in the guard cells of WT poplar and transgenic lines. (d) Detection of ABA-induced stomatal closure in the leaves of the WT and transgenic lines; leaves were separated and immersed under light in stomata-opening solution (OS) for 2 h and then treated with 5 μ M ABA for 2 hours (OS \rightarrow ABA). (e) Stomatal closure was observed at 0, 1 and 2 h of ABA treatment with scanning electron microscopy of stomatal aperture. Scale bars = 50 μ m. Error bars are means \pm SE (n = 50). Asterisks denote significant differences: *, P \leq 0.05; **, P \leq 0.01.

Figure 5. Light response curves were measured in WT, *OXPeCHYR1-1*, and *OXPeCHYR1-8*.

Light response curves were measured in the same greenhouse conditions. (a) A-light curve. (b) Gs-light curve. (c) Transpiration-light curve. (d) Instantaneous WUE-light curve. (e) VPD-light curve. Data are means \pm SE (n = 25). Asterisks denote significant differences: *, P \leq 0.05; **, P \leq 0.01.

Figure 6. *35S:PeCHYRI* plants exhibited increased tolerance during short-term drought treatment. (a) Morphological differences in short-term drought assays. Bar = 5 cm. Quantitative measurement of leaf RWC (b), MDA content analysis (c), REC (relative electrical conductance) (d), H₂O₂ (e), activity of peroxidase (POD) (f), and activity of superoxide dismutase (SOD) (g) in the leaves of WT and *35S:PeCHYRI* plants under normal and drought stress conditions. Values are means \pm SE (n = 40). All asterisks denote significant differences: *, P \leq 0.05; **, P \leq 0.01.

Figure 7. Variation in photosynthetic parameters of *35S:PeCHYRI* plants relative to those of WT plants during drought treatment and changes in the water loss rate of *35S:PeCHYRI* plants relative to that of WT plants during natural dehydration. (a) Transpiration-drought time curve. (b) Gs-drought time curve. (c) A-drought time curve. (d) Water loss from detached leaves is indicated as a percentage of initial fresh weight. Values are means \pm SE (n = 25). All asterisks denote significant differences: *, P \leq 0.05; **, P \leq 0.01.

Figure 8. *35S:PeCHYRI* poplars exhibited high drought tolerance during long-term drought. (a) Morphological differences between WT and transgenic plants during long-term drought assays. Bar = 6 cm. The contents of chlorophyll a and b (b), the maximum photosynthetic efficiency of photosystem II (Fv/Fm) (c), and leaf relative water content (RWC) (d) under 70% soil RWC and 45% soil RWC. Data are means \pm SE (n = 40). Asterisks denote significant differences: *, P \leq 0.05; **, P \leq 0.01.

Figure 9. *PeCHYRI* positively regulates plant height and biomass increase under long-term drought. (a) Plant height under 70% soil RWC and 45% soil RWC. (b) Stem height growth rate under 70% soil RWC and 45% soil RWC. (c) Shoot biomass, (d) root biomass, (e) whole plant biomass, and (f) root-shoot ratio in WT and *35S:PeCHYRI* poplars under no

water stress and mild water stress. Data are means \pm SE (n = 25). Asterisks denote significant differences: *, $P \leq 0.05$; **, $P \leq 0.01$.

Figure 10. Stress-responsive genes expressed in WT and 35S:*PeCHYR1* poplar plants under non-stress conditions and instantaneous drought stress conditions. Relative expression levels of *PeSnrk2.6* (A), *PeLea14* (B), *PeRbohD* (C), and *PeRbohF* (D) in WT and 35S:*PeCHYR1* poplar plants under non-stress conditions and 2 h drought stress. Total RNA was extracted from 2-week-old seedlings treated with drought stress for 2 h, and RT-qPCR was performed. Data are means \pm SE (n = 20). Asterisks denote significant differences: *, $P \leq 0.05$; **, $P \leq 0.01$.

Supplement materials

Figure S1. Analysis of Cis elements of the *PeCHYR1* gene.

Figure S2. Analysis of the transgenic poplar plants overexpressing *PeCHYR1*.

Figure S3. 35S: *PeCHYR1* plants exhibited increased disease resistance under anthracnose fungus treatment.

Table S1. Primer sequences used for cloning of *PeCHYR1* cDNA and RT-PCR.

Table S2. Accession numbers of gene.









